

**CANDIX
MANUFACTURER**
<http://psresearch.com/product>

COMPOSITION
Clindamycin (2% w/w) + Clotrimazole (2% w/w)

STORAGE
Store below 30°C



shutterstock.com - 115276195

shutterstock.com - 115276195

PRODUCT INTRODUCTION

Candid-CL Vaginal gel is a combination medicine that is prescribed to treat various types of vaginal infections associated with vaginal discharge. It fights against the infection by stopping the growth of infection-causing microorganisms. It also prevents the further spread of the infection.

Candid-CL Vaginal gel is to be used as per the label instructions or as suggested by your doctor. This medicine is only for vaginal application. You must wash your hands before and after using it. It is best to use this at bedtime regularly to get the most benefit. If there is any accidental exposure of the medicine to the eyes or mouth, you must wash it immediately. For better efficacy of the medicine, the course of the treatment should be completed even if you feel better. Do not use it more than the recommended dose.

Candid-CL Vaginal gel may lead to a few common side effects like mild irritation and burning sensation at the application site. If these side effects do not improve with time or worsen, inform your doctor. In case of any allergic reactions (rashes, itching, swelling, shortness of breath, etc.), it is advised to call for immediate medical help.

Before using it, tell your doctor if you have an underlying disease or are taking medicines for any health condition. The medicines that are used externally usually do not get affected by other medicines, however, you should let your doctor know if you are taking any medicines to ensure safety. Ask your doctor for advice if you are pregnant or breastfeeding. If you are a known allergic to the medicine, you must refrain from using this medicine and consult your doctor. During the course of the treatment, you must follow proper personal hygiene for a faster recovery.

USES OF CANDID-CL VAGINAL GEL

- Syndromic treatment of vaginal discharge

BENEFITS OF CANDID-CL VAGINAL GEL

In Syndromic treatment of vaginal discharge

Vaginal discharge is completely normal if it is not accompanied by other symptoms such as foul smell, change in color, sudden increase in the quantity, itching, irritation or burning sensation. These symptoms can be embarrassing as well as bothersome. Candid-CL Vaginal gel helps in effectively managing these symptoms by killing the organisms responsible for causing an infection of the vagina that can lead to such offensive symptoms.

Take it as prescribed by the doctor to get maximum benefit. Besides, always maintain good personal hygiene to avoid any vaginal infection. Always wipe from front to back after using the toilet each time, wear cotton underpants, bathe daily and avoid use of any sprays or deodorised products in the vaginal area. Consult your doctor if you have any doubts.

SIDE EFFECTS OF CANDID-CL VAGINAL GEL

Most side effects do not require any medical attention and disappear as your body adjusts to the medicine. Consult your doctor if they persist or if you're worried about them

Common side effects of Candid-CL

- Vaginal burning sensation
- Itching
- Irritation

HOW TO USE CANDID-CL VAGINAL GEL

This medicine is for external use only. Use it in the dose and duration as advised by your doctor. Check the label for directions before use. Clean and dry the affected area and apply the gel. Wash your hands after applying

HOW CANDID-CL VAGINAL GEL WORKS

Candid-CL Vaginal gel is a combination of two medicines: Clindamycin and Clotrimazole. Clindamycin is an antibiotic which stops bacterial growth in the vaginal tract by preventing the synthesis of essential proteins required by bacteria to carry out vital functions. Clotrimazole is an antifungal which stops the growth of fungi by preventing them from making their protective covering. Together, they treat your vaginal infection effectively.

SAFETY ADVICE

Alcohol

No interaction found/established

Pregnancy

SAFE IF PRESCRIBED

Candid-CL Vaginal gel is generally considered safe to use during pregnancy. Animal studies have shown low or no adverse effects to the developing baby; however, there are limited human studies.

Breast feeding

SAFE IF PRESCRIBED

Candid-CL Vaginal gel is safe to use during breastfeeding. Human studies suggest that the drug does not pass into the breastmilk in a significant amount and is not harmful to the baby.

Driving

No interaction found/established

Kidney

No interaction found/established

Liver

No interaction found/established

WHAT IF YOU FORGET TO TAKE CANDID-CL VAGINAL GEL?

If you miss a dose of Candid-CL Vaginal gel, use it as soon as possible. However, if it is almost time for your next dose, skip the missed dose and go back to your regular schedule. Do not double the dose.

All substitutes

For informational purposes only. Consult a doctor before taking any medicines.

Candid-CL Vaginal gel is inserted into the vagina using an applicator, usually at night before going to bed.

- Do not skip any doses and finish the full course of treatment even if you feel better.
 - Avoid contact with your eyes. If accidental exposure occurs, rinse immediately with plenty of water.
 - It may cause minor burning, stinging, or irritation when applied. Inform your doctor if this does not go away.
 - It can damage condoms and diaphragms and make them ineffective at preventing pregnancy. Ask your doctor about other suitable methods of preventing pregnancy.
 - Inform your doctor if your symptoms do not get improved within 3 days of taking the medicine.
-